



7<sup>th</sup> EFSMA - European Sports Medicine Congress

3<sup>rd</sup> Central European Congress of Physical Medicine and Rehabilitation

Annual Assembly of the German and the Austrian Society of Physical Medicine and Rehabilitation

Salzburg, 26th - 29th October 2011



# 7<sup>th</sup> EFSMA – European Congress of Sports Medicine

## 3<sup>rd</sup> Central European Congress of Physical Medicine and Rehabilitation

### Annual Assembly of the German and the Austrian Society of Physical Medicine and Rehabilitation

Austrian Society of Physical Medicine and Rehabilitation

Salzburg, Austria  
26<sup>th</sup>–29<sup>th</sup> October 2011

Book of Abstracts



This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 License.

<http://creativecommons.org/licenses/by-nc-nd/3.0/deed.en>

Organizer: ÖGPMR – Österreichische Gesellschaft für Physikalische Medizin und Rehabilitation

President: Prof. Anton Wicker

Layout: Editorial Office German Medical Science

The abstracts are published online on the German Medical Science website

<http://www.egms.de/en/meetings/esm2011/>

## Physical activity, isometric strength and body composition of university students

Leonel Preto, Eugénia Mendes, André Novo, Ana Azevedo

Polytechnic Institute of Bragança – School of Health, Bragança, Portugal

**Objective:** The importance of physical activity in promoting health and well-being of the young adult population is evidenced by several epidemiological studies. The increasing incidence in this population of cardiovascular, metabolic and degenerative diseases and obesity has been linked with an epidemic variable, the sedentary behavior, and commonly dubbed the “disease of the century”. Regular physical activity may contribute to the development of muscle strength, improve the quality of soft tissue and bone mass, and decreased body fat. The early inclusion of regular physical activity and other healthy lifestyles tend to be incorporated throughout life with noticeable gains in health and quality of life in adult life.

In this study we assessed the physical activity level (PAL), muscle strength (MS) and body composition (BC) of 84 young adults in higher education with the following objectives: assess physical activity; evaluate different manifestations of muscle strength; assess body composition and analyze the relationship between muscle strength, body composition and physical activity in young adults.

**Material/Methods:** We drew up a descriptive, correlational and cross-cutting study. To achieve our objective we collected the following data:

- a) Physical activity: short version of IPAQ (International Physical Activity Questionnaire).
- b) Muscle strength: hand grip of both hands with Jamar® Hand Dynamometer;
- c) Key pinch strength with digital dynamometer Baseline®;
- d) Maximum isometric strength of the quadriceps in seat 45-degree incline leg press machine linked to a strain gauge Ergo Meter – Globus® and appropriated software
- f) Body composition: bioelectric impedance on Tanita Ironman Body Composition Monitor®
- g) Height: stadiometer Seca®

**Results:** A total of 84 university students were subject to this study, 66 females and 18 males with an average of  $20,9 \pm 2,3$  years old. In IPAQ classification we found 50 students with low, 25 with moderate and 9 with high level of physical activity. Relatively to muscle strength there were significant correlations between handgrip strength (both hands), key pinch strength (both hands) and isometric maximum strength of the quadriceps. Analysing body composition we can observe that female students with more IMC and more body fat present higher levels of physical activity. Regarding the male students, we can find the opposite: the lesser the percentage of body fat and lower IMC levels the higher physical activity. We found no relation between the different strength demonstrations and the levels of physical activity. When correlating handgrip strength (right hand; left hand) with body composition, we observed correlations between:

- Height (0,704\*\*; 0,705\*\*), total body fat (-0,412\*\*; -0,428\*\*), weight (0,406\*\*; 0,463\*\*), body water (0,441\*\*; 0,453\*\*), visceral fat level (0,223\*; 0,259\*), bone mass (0,800\*\*; 0,798\*\*) and muscular mass (0,816\*\*; 0,803\*\*).

When correlating key pinch strength (right hand; left hand) with body composition, we observed correlations between:

- Height (0,577\*\*; 0,531\*\*), total body fat (-0,389\*\*; -0,252\*), weight (0,336\*\*; 0,424\*\*), body water (0,449\*\*; 0,348\*\*), bone mass (0,638\*\*; 0,657\*\*) and muscular mass (0,677\*\*; 0,684\*\*).

When correlating maximum isometric strength of the quadriceps with body composition, we observed correlations between:

- Height (0,355\*\*), total body fat (-0,389\*\*), body water (0,389\*\*), bone mass (0,480\*\*) and muscular mass (0,490\*\*).

**Conclusion:** Results indicate that the evaluated university students are mainly sedentary. We also concluded that there are very strong positive correlations between the different manifestations of strength and that there is no relationship between the different manifestations of strength and levels of physical activity. Analyzing our results, we can conjecture about why young people do exercise: females exercise themselves to lose weight and males to maintain physical condition. But there are strong correlations between the different manifestations of strength and the different variables of body composition.

## References

1. Fogelholm M. Physical activity, fitness and fatness: relations to mortality, morbidity and disease risk factors. A systematic review. *Obes Rev.* 2010;11(3):202-21.
2. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc.* 2007;39(8):1423-34.
3. Lafortuna CL, Maffiuletti NA, Agosti F, Sartorio A. Gender variations of body composition, muscle strength and power output in morbid obesity. *Int J Obes (Lond).* 2005;29(7):833-41.
4. Peters MJ, van Nes SI, Vanhoutte EK, Bakkers M, van Doorn PA, Merckies IS, et al. Revised normative values for grip strength with the Jamar dynamometer. *J Peripher Nerv Syst.* 2011;16(1):47-50.
5. Thompson AM, Humbert ML, Mirwald RL. A longitudinal study of the impact of childhood and adolescent physical activity experiences on adult physical activity perceptions and behaviors. *Qual Health Res.* 2003;13(3):358-77.

Please cite as: Preto L, Mendes E, Novo A, Azevedo A. Physical activity, isometric strength and body composition of university students. In: 7th EFSMA – European Congress of Sports Medicine, 3rd Central European Congress of Physical Medicine and Rehabilitation. Salzburg, 26.-29.10.2011. Düsseldorf: German Medical Science GMS Publishing House; 2011. Doc11esm229.

DOI: 10.3205/11esm229, URN: urn:nbn:de:0183-11esm2295

Freely available from: <http://www.egms.de/en/meetings/esm2011/11esm229.shtml>